

Index

A

- Abadir, Karim M. v
- added variable plot 22
- adjoint matrix 87
- admissibility 69
- Albert's theorem 114
- ANOVA 41
 - another parametrization 43
 - conditional mean and variance 42
 - estimability 52
 - in matrix terms 42
- antieigenvalues 62, 65
- approximating
 - the centered data matrix 79
- AR(1)-structure *see* autocorrelation
- autocorrelation
 - $\mathbf{V} = \{\rho^{|i-j|}\}$ 34
 - BLP 35
 - in linear model 36
 - in prediction 71
 - regression coefficients 36
 - testing 36

B

- Baksalary, Oskar Maria vi
- Banachiewicz–Schur form 93
- Bernoulli distribution 23
- Bernstein, Dennis S. vi
- best linear predictor *see* BLP
- best linear unbiased estimator *see* BLUE
- best linear unbiased predictor *see* BLUP, 69
- best predictor *see* BP
- bias 31
- binomial distribution 23
- Blanck, Alice vi

- block diagonalization 32, 93
- Bloomfield–Watson efficiency 66
- Bloomfield–Watson–Knott inequality 62
- BLP
 - BLP($\mathbf{y}; \mathbf{x}$) 26, 32
 - BLP($\mathbf{z}; \mathbf{A}'\mathbf{z}$) 33
 - BLP($y_n; \mathbf{y}_{(n-1)}$) 35
 - $\boldsymbol{\mu}_y + \boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1} (\mathbf{x} - \boldsymbol{\mu}_x)$ 26
 - $\boldsymbol{\mu}_y + \boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1} (\mathbf{x} - \boldsymbol{\mu}_x)$ 32
 - autocorrelation 35
- BLUE
 - $\tilde{\boldsymbol{\beta}}$, representations 54
 - $\tilde{\boldsymbol{\beta}}_2 = (\mathbf{X}'_2 \dot{\mathbf{M}}_{1w} \mathbf{X}_2)^{-1} \mathbf{X}'_2 \dot{\mathbf{M}}_{1w} \mathbf{y}$ 57
 - $\tilde{\boldsymbol{\beta}}_2 = (\mathbf{X}'_2 \dot{\mathbf{M}}_1 \mathbf{X}_2)^{-1} \mathbf{X}'_2 \dot{\mathbf{M}}_1 \mathbf{y}$ 57
 - $\tilde{\boldsymbol{\beta}}_i$ in the general case 57
 - BLUE($\mathbf{K}'\boldsymbol{\beta}$) 53
 - BLUE($\mathbf{X}\boldsymbol{\beta}$) 52
 - definition 52
 - fundamental equation 53
 - cov($\mathbf{X}\tilde{\boldsymbol{\beta}}$) as a shorted matrix 115
 - $\mathbf{X}\tilde{\boldsymbol{\beta}}$, general representations 53
 - $\mathbf{X}\tilde{\boldsymbol{\beta}}$, representations 54
 - equality of OLSE and BLUE, several conditions 58
 - restricted BLUE 41
 - without the i th observation 47
- BLUP
 - fundamental equation 70
 - in mixed model 72
 - of y_* 18
 - of \boldsymbol{y} 72
 - of \mathbf{y}_f 70
 - Pandora's Box 70
 - representations for BLUP(\mathbf{y}_f) 70
- BP
 - BP($\mathbf{y}; \mathbf{x}$) 27

- $BP(y; \mathbf{x})$ 31
 $E(y | \mathbf{x})$ 31
 in N_p 27
 is the conditional expectation 27, 31
- C**
- canonical correlations
 between \mathbf{x} and \mathbf{y} 80
 between $\hat{\boldsymbol{\beta}}$ and $\tilde{\boldsymbol{\beta}}$ 63
 between $\mathbf{H}\mathbf{y}$ and $\mathbf{M}\mathbf{y}$ 64
 between $\mathbf{K}'\mathbf{Q}_L\mathbf{u}$ and $\mathbf{L}'\mathbf{Q}_K\mathbf{u}$ 82
 non-unit cc's between $\mathbf{K}'\mathbf{u}$ and $\mathbf{L}'\mathbf{u}$ 82
 proper eigenvalues 81
 unit cc's 64
 Watson efficiency 63
 Cauchy–Schwarz inequality 116
 centered $\mathbf{X}_0, \mathbf{X}_y$ 6
 centered and scaled \mathbf{X}_y 7
 centered model 15
 centering matrix \mathbf{C} 6
 characteristic equation 100
 Chatterjee, Samprit v
 Cholesky decomposition 111
 Cochran's theorem 28
 coefficient of determination *see* multiple correlation
 cofactor 86
 collinearity 46
 column space
 definition 9
 $\mathcal{C}(\mathbf{V}\mathbf{X}^\perp)$ 99
 $\mathcal{C}(\mathbf{W} - \mathbf{X} : \mathbf{I} - \mathbf{W}\mathbf{W})$ 99
 $\mathcal{C}(\mathbf{X} : \mathbf{V}) = \mathcal{C}(\mathbf{X} : \mathbf{V}\mathbf{M})$ 53
 $\mathcal{C}[\mathbf{X}'(\mathbf{V} + \mathbf{X}\mathbf{U}\mathbf{X}')^{-1}\mathbf{X}]$ 50
 simple properties 82
 of $\text{cov}(\mathbf{X}\tilde{\boldsymbol{\beta}})$ 55
 completely symmetric matrix 61
 eigenvalues 101, 102
 condition index 45
 condition number 45
 conditional *see* normal distribution
 $E[E(y | \mathbf{x})]$ 27
 $\text{var}[E(y | \mathbf{x})]$ 27, 42
 expectation and the best predictor 31
 confidence ellipse *see* confidence region
 confidence interval
 for β_i 18
 for $E(y_*)$ 19
 Working–Hotelling confidence band 19
 confidence region
 for $\boldsymbol{\beta}_2$ 38
 for $\boldsymbol{\beta}_x$ 38
 for $\boldsymbol{\mu}$ 29
- consistency (when solving \mathbf{X})
 of $\mathbf{A}\mathbf{X} = \mathbf{B}$ 91
 of $\mathbf{A}\mathbf{X}\mathbf{B} = \mathbf{C}$ 91
 of $\mathbf{X}(\mathbf{A} : \mathbf{B}) = (\mathbf{0} : \mathbf{B})$ 85, 98
 consistency of linear model 53
 constraints to obtain unique $\hat{\boldsymbol{\beta}}$ 39
 contingency table 23
 contours of constant density 24, 121
 contrast 52
 Cook's distance 45
 $\text{COOK}_i^2(V)$ 47
 Cook, R. Dennis 34
 correlation *see* multiple correlation
 $\text{cor}(x, y) = \sigma_{xy}/\sigma_x\sigma_y$ 3
 $\text{cor}_s(x_i, x_j), \text{cor}_d(\mathbf{x}_i, \mathbf{x}_j)$ 8
 $\mathbf{R} = \text{cor}_d(\mathbf{y}, \hat{\mathbf{y}})$ 14
 $\text{cor}(\hat{\boldsymbol{\beta}}_1, \hat{\boldsymbol{\beta}}_2)$ 18, 22
 $\text{cor}(\hat{\boldsymbol{\epsilon}}_i, \hat{\boldsymbol{\epsilon}}_j)$ 14
 as a cosine 8
 between dichotomous variables 23
 random sample without replacement 23
 correlation matrix
 $\text{cor}(\mathbf{x})$ 10
 $\text{cor}_d(\mathbf{X}_0) = \mathbf{R}_{xx}$ 7
 $\text{cor}_d(\mathbf{X}_0 : \mathbf{y}) = \mathbf{R}$ 7
 “correlation matrix” 114
 determinant 88, 89
 inverse 17, 89, 90
 rank 9, 88
 recursive decomposition of $\det(\mathbf{R})$ 89
 Courant–Fischer theorem 104
 covariance
 $\text{cov}(x, y) = E(x - \mu_x)(y - \mu_y)$ 3
 $\text{cov}_s(x, y), \text{cov}_d(\mathbf{x}, \mathbf{y})$ 8
 $\text{cov}(\mathbf{z}'\mathbf{A}\mathbf{z}, \mathbf{z}'\mathbf{B}\mathbf{z})$ 28
 covariance matrix
 $\text{cov}(\mathbf{x})$ 10
 $\text{cov}(\mathbf{x}, \mathbf{y})$ 10
 $\text{cov}(\hat{\boldsymbol{\beta}})$ 16
 $\text{cov}(\hat{\boldsymbol{\beta}}_2)$ 17
 $\text{cov}(\hat{\boldsymbol{\beta}}_x, \bar{y}) = \mathbf{0}$ 18
 $\text{cov}(\tilde{\boldsymbol{\beta}}_2)$ 57
 $\text{cov}(\mathbf{e}_{y-x})$ 32
 $\text{cov}(\mathbf{e}_{y-x}, \mathbf{x})$ 32
 $\text{cov}(\mathbf{X}\tilde{\boldsymbol{\beta}})$ 55
 $\text{cov}(\mathbf{x}, \mathbf{y} - \mathbf{A}\mathbf{x}) = \mathbf{0}$ 33
 $\text{cov}[\mathbf{z} - \text{BLP}(\mathbf{z}; \mathbf{A}'\mathbf{z})]$ 33
 $\text{cov}_d(\mathbf{U})$ 11
 $\text{cov}_d(\mathbf{U}\mathbf{A})$ 11
 $\text{cov}_d(\mathbf{U}\mathbf{S}^{-1/2})$ 11
 $\text{cov}_d(\mathbf{U}\mathbf{T}), \mathbf{S} = \mathbf{T}\mathbf{A}\mathbf{T}'$ 11
 $\text{cov}_d(\mathbf{X}_0) = \mathbf{S}_{xx}$ 7
 $\text{cov}_d(\mathbf{X}_y) = \mathbf{S}$ 7

always nonnegative definite 10
 of the prediction error 32, 78
 partial covariance 26
 crazy, still ν
 Cronbach's α 118

D

data matrix
 $\mathbf{X}_y = (\mathbf{X}_0 : \mathbf{y})$ 6
 best rank- k approximation 79
 keeping data as a theoretical distribution 8
 decomposition *see* eigenvalue decomposition, *see* full rank decomposition, *see* singular value decomposition
 Hartwig–Spindelböck 112
 derivatives, matrix derivatives 120
 determinant
 $\Sigma_{2 \times 2}$ 25
 Laplace expansion 86
 recursive decomposition of $\det(\mathbf{S})$ 34
 recursive decomposition of $\det(\Sigma)$ 34
 Schur complement 87
 DFBETA $_i$ 45
 diagonalizability 103
 discrete uniform distribution 23
 discriminant function 79
 disjointness
 $\mathcal{C}(\mathbf{A}) \cap \mathcal{C}(\mathbf{B}) = \{\mathbf{0}\}$, several conditions 84
 $\mathcal{C}(\mathbf{VH}) \cap \mathcal{C}(\mathbf{VM})$ 64
 $\mathcal{C}(\mathbf{X}_{(1)}) \cap \mathcal{C}(\mathbf{X}_{(2)})$ 47
 $\mathcal{C}(\mathbf{X}) \cap \mathcal{C}(\mathbf{VM})$ 53
 $\mathcal{C}(\mathbf{X}_1) \cap \mathcal{C}(\mathbf{X}_2)$ 12
 estimability in outlier testing 46
 distance *see* Cook's distance, *see* Mahalanobis distance, *see* statistical distance
 distribution *see* Hotelling's T^2 , *see* normal distribution
 F -distribution 28
 t -distribution 28
 Bernoulli 23
 binomial 23
 chi-squared 27, 28
 discrete uniform 23
 Hotelling's T^2 29
 of $\text{MHLN}^2(\mathbf{x}, \boldsymbol{\mu}, \Sigma)$ 27
 of $\mathbf{y}'(\mathbf{H} - \mathbf{J})\mathbf{y}$ 28
 of $\mathbf{y}'(\mathbf{I} - \mathbf{H})\mathbf{y}$ 28
 of $\mathbf{z}'\mathbf{A}\mathbf{z}$ 27, 28
 of $\mathbf{z}'\Sigma^{-1}\mathbf{z}$ 27
 Wishart-distribution 28, 77
 dominant eigenvalue 101, 112

Draper, Norman R. ν
 Durbin–Watson test statistic 36
 Dürer's *Melencolia I* 113

E

Eckart–Young theorem
 non-square matrix 111
 symmetric matrix 105
 efficiency of OLSE *see* Watson efficiency, Bloomfield–Watson efficiency, Rao's efficiency
 eigenvalue, dominant 101, 112
 eigenvalues
 definition 100
 (λ, \mathbf{w}) as an eigenpair for (\mathbf{A}, \mathbf{B}) 106
 $\text{ch}(\Sigma_{2 \times 2})$ 25
 $\text{ch}(\mathbf{A}, \mathbf{B}) = \text{ch}(\mathbf{B}^{-1}\mathbf{A})$ 106
 $\text{ch}(\mathbf{A}_{2 \times 2})$ 100
 $\text{ch}_1(\mathbf{A}, \mathbf{B}) = \text{ch}_1(\mathbf{B}^+\mathbf{A})$ 108
 $|\mathbf{A} - \lambda\mathbf{I}| = \mathbf{0}$ 100
 $|\mathbf{A} - \lambda\mathbf{B}| = \mathbf{0}$ 105
 $\text{nzch}(\mathbf{A}, \mathbf{B}) = \text{nzch}(\mathbf{A}\mathbf{B}^-)$ 108
 $\text{nzch}(\mathbf{A}\mathbf{B}) = \text{nzch}(\mathbf{B}\mathbf{A})$ 103
 algebraic multiplicity 103
 characteristic polynomial 100
 eigenspace 103
 geometric multiplicity 103
 intraclass correlation 101, 102
 of $\alpha^2\mathbf{I} + \mathbf{a}\mathbf{1}' + \mathbf{1}\mathbf{a}'$ 105
 of $\text{cov}(\mathbf{X}\hat{\boldsymbol{\beta}})$ 108
 of $\mathbf{P}_A\mathbf{P}_B$ 82, 96, 97
 of $\mathbf{P}_K\mathbf{P}_L - \mathbf{P}_F$ 82
 spectral radius 101
 elementary column/row operation 103
 equality
 of BLUPs of \mathbf{y}_f under two models 71
 of BLUPs under two mixed models 74
 of OLSE and BLUE, several conditions 58
 of OLSE and BLUE, subject to
 $\mathcal{C}(\mathbf{U}) \subset \mathcal{C}(\mathbf{X})$ 60
 of the BLUEs under two models 59
 of the OLSE and BLUE of $\boldsymbol{\beta}_2$ 61
 $\hat{\boldsymbol{\beta}}(\mathcal{M}_{(i)}) = \tilde{\boldsymbol{\beta}}(\mathcal{M}_{(i)})$ 66
 $\hat{\boldsymbol{\beta}}_2(\mathcal{M}_{12}) = \tilde{\boldsymbol{\beta}}_2(\mathcal{M}_{12})$ 61
 $\tilde{\boldsymbol{\beta}}_2(\mathcal{M}_{12}) = \hat{\boldsymbol{\beta}}_2(\mathcal{M}_{12})$ 67
 BLUP(\mathbf{y}_f) = BLUE($\mathbf{X}_f\boldsymbol{\beta}$) 70
 BLUP(\mathbf{y}_f) = OLSE($\mathbf{X}_f\boldsymbol{\beta}$) 70
 $\text{SSE}(V) = \text{SSE}(I)$ 57
 $\mathbf{X}\hat{\boldsymbol{\beta}} = \mathbf{X}\tilde{\boldsymbol{\beta}}$ 58
 $\mathbf{A}\mathbf{y} = \mathbf{B}\mathbf{y}$ 54
 $\mathbf{P}_{\mathbf{X};v_1} = \mathbf{P}_{\mathbf{X};v_2}$ 59
 estimability

definition 51
 constraints 39
 contrast 52
 in hypothesis testing 37, 41
 of $\boldsymbol{\beta}$ 52
 of β_k 52
 of $\boldsymbol{\delta}$ in the extended model 46
 of δ in the extended model 44
 of $\mathbf{K}'\boldsymbol{\beta}$ 51
 of $\mathbf{X}_2\boldsymbol{\beta}_2$ 51
 of $\mathbf{c}'\boldsymbol{\tau}$ 52
 extended (mean-shift) model 44, 46, 47

F

factor analysis 79
 factor scores 80
 finiteness matters 24
 fitted values 12
 frequency table 23
 Frisch–Waugh–Lovell theorem 15, 22
 generalized 61
 Frobenius inequality 84
 full rank decomposition 92, 111

G

Galton's discovery of regression v
 Gauss–Markov model *see* linear model
 Gauss–Markov theorem 53
 generalized inverse
 the 4 Moore–Penrose conditions 90
 Banachiewicz–Schur form 93
 least-squares g-inverse 94
 minimum norm g-inverse 94
 partitioned nnd matrix 93
 reflexive g-inverse 91, 93
 through SVD 92
 generalized variance 10, 11
 generated observations from $N_2(\mathbf{0}, \boldsymbol{\Sigma})$ 121

H

Hadi, Ali S. v
 Hartwig–Spindelböck decomposition 112
 Haslett, Stephen J. vi
 hat matrix \mathbf{H} 12
 Henderson's mixed model equations 73
 Hotelling's T^2 28, 29, 77
 when $n_1 = 1$ 30
 hypothesis
 $\boldsymbol{\beta}_2 = \mathbf{0}$ 38
 $\boldsymbol{\beta}_x = \mathbf{0}$ 31, 40
 $\boldsymbol{\delta} = \mathbf{0}$ in outlier testing 46

$\boldsymbol{\mu} = \boldsymbol{\mu}_0$ 29
 $\boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$ 30, 77
 $\boldsymbol{\beta}_1 = \cdots = \boldsymbol{\beta}_k = \mathbf{0}$ 40
 $\beta_i = 0$ 40
 $\mathbf{k}'\boldsymbol{\beta} = 0$ 40
 $\boldsymbol{\delta} = \mathbf{0}$ in outlier testing 44
 $\mu_1 = \cdots = \mu_g$ 41
 $\mu_1 = \mu_2$ 42
 $\mathbf{K}'\boldsymbol{\beta} = \mathbf{d}$ 39, 40
 $\mathbf{K}'\boldsymbol{\beta} = \mathbf{d}$ when $\text{cov}(\mathbf{y}) = \sigma^2\mathbf{V}$ 40
 $\mathbf{K}'\boldsymbol{\beta} = \mathbf{0}$ 37
 $\mathbf{K}'\mathbb{B} = \mathbf{D}$ 77
 overall- F -value 40
 testable 41
 under intraclass correlation 59

I

idempotent matrix
 eigenvalues 97
 full rank decomposition 97
 rank = trace 97
 several conditions 97
 illustration of $\text{SST} = \text{SSR} + \text{SSE}$ 121
 independence
 linear 9
 statistical 24
 $\bar{\mathbf{u}}$ and $\mathbf{T} = \mathbf{U}'(\mathbf{I} - \mathbf{J})\mathbf{U}$ 29
 observations, rows 28, 29
 of dichotomous variables 24
 quadratic forms 27, 28
 inequality
 $\text{ch}_i(\mathbf{A}\mathbf{A}') \leq \text{ch}_i(\mathbf{A}\mathbf{A}' + \mathbf{B}\mathbf{B}')$ 103
 $\lambda_n \leq \mathbf{x}'\mathbf{A}\mathbf{x}/\mathbf{x}'\mathbf{x} \leq \lambda_1$ 104
 $\|\mathbf{A}\mathbf{x}\| \leq \|\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}\|$ 93
 $\|\mathbf{A}\mathbf{x}\|_2 \leq \|\mathbf{A}\|_2 \|\mathbf{x}\|_2$ 110
 $\text{tr}(\mathbf{A}\mathbf{B}) \leq \text{ch}_1(\mathbf{A}) \cdot \text{tr}(\mathbf{B})$ 103
 Bloomfield–Watson–Knott 62
 Cauchy–Schwarz 116
 Frobenius 84
 Kantorovich 116
 Samuelson's 118
 Sylvester's 84
 Wielandt 117
 inertia 115
 intercept 14
 interlacing theorem 103
 intersection
 $\mathcal{C}(\mathbf{A}) \cap \mathcal{C}(\mathbf{B})$ 84
 $\mathcal{C}(\mathbf{A}) \cap \mathcal{C}(\mathbf{B})$, several properties 96
 intraclass correlation
 F -test 59
 OLSE = BLUE 59
 eigenvalues 101, 102

invariance
 of $\mathcal{C}(\mathbf{AB}^{-1}\mathbf{C})$ 91
 of $\mathbf{AB}^{-1}\mathbf{C}$ 91
 of $\mathbf{X}'(\mathbf{V} + \mathbf{XUX}')^{-1}\mathbf{X}$ 50
 of $r(\mathbf{AB}^{-1}\mathbf{C})$ 91
 inverse
 of $\mathbf{A}_{n \times n}$ 86
 intraclass correlation 101
 of $\mathbf{A} = \{a_{ij}\} = \{\min(i, j)\}$ 36
 of \mathbf{R} 89
 of \mathbf{R}_{xx} 17, 90
 of $\mathbf{X}'\mathbf{X}$ 16, 88
 of $\Sigma_{2 \times 2}$ 25
 of a partitioned block matrix 87
 of autocorrelation matrix 35
 of partitioned pd matrix 88
 Schur complement 87
 irreducible matrix 112

K

Kantorovich inequality 116
 Kronecker product 119
 in multivariate model 76
 in multivariate sample 29

L

Lagrangian multiplier 39
 Laplace expansion of determinant 86
 $\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$ vi
 least squares *see* OLSE
 Lee, Alan J. v
 Lehmann–Scheffé theorem 68
 leverage h_{ii} 45
 likelihood function 30
 likelihood ratio 30
 linear hypothesis *see* hypothesis
 linear independence 9
 collinearity 46
 linear model 5
 multivariate 76
 linear prediction sufficiency 72
 linear zero function 54, 70
 linearly complete 68
 linearly sufficient *see* sufficiency
 Löwner ordering *see* minimizing
 definition 113
 Albert's theorem 114
 $\mathbf{Y}'\mathbf{M}\mathbf{Y} \leq_L (\mathbf{Y} - \mathbf{XB})'(\mathbf{Y} - \mathbf{XB})$ 22, 117

M

magic square 113

Magnus, Jan R. v
 Mahalanobis distance
 $\text{MHLN}^2(\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \mathbf{S}_*)$ 30, 77, 79
 $\text{MHLN}^2(\mathbf{u}_{(i)}, \bar{\mathbf{u}}, \mathbf{S})$ 8
 $\text{MHLN}^2(\mathbf{x}, \boldsymbol{\mu}, \boldsymbol{\Sigma})$ 9, 27
 $\text{MHLN}^2(\mathbf{x}_*, \bar{\mathbf{x}}, \mathbf{S}_{xx})$ 18
 $\text{MHLN}^2(\mathbf{x}_{(i)}, \bar{\mathbf{x}}, \mathbf{S}_{xx})$ 45, 118
 $\text{MHLN}^2(\boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ 79
 major/minor axis of the ellipse 25
 matrix
 irreducible 112
 nonnegative 112
 permutation 112
 reducible 112
 shorted 115
 spanning $\mathcal{C}(\mathbf{A}) \cap \mathcal{C}(\mathbf{B})$ 84
 stochastic 113

matrix $\ddot{\mathbf{M}}$
 $\mathbf{P}_V \ddot{\mathbf{M}} \mathbf{P}_V$ 48
 $\mathbf{P}_W \ddot{\mathbf{M}} \mathbf{P}_W = \ddot{\mathbf{M}}_W$ 49

matrix $\dot{\mathbf{M}}$
 $\mathbf{M}(\mathbf{M}\mathbf{V}\mathbf{M})^{-1}\mathbf{M}$ 48
 $\text{SSE}(\mathbf{Y}) = \mathbf{y}'\dot{\mathbf{M}}\mathbf{y}$ 56, 57

matrix $\dot{\mathbf{M}}_1$
 $\mathbf{M}_1(\mathbf{M}_1\mathbf{V}\mathbf{M}_1)^{-1}\mathbf{M}_1$ 57, 99

matrix $\dot{\mathbf{M}}_{1W}$
 $\mathbf{M}_1(\mathbf{M}_1\mathbf{W}_2\mathbf{M}_1)^{-1}\mathbf{M}_1$ 57

matrix angle 65
 matrix of corrected sums of squares
 $\mathbf{T} = \mathbf{X}'_y \mathbf{C}\mathbf{X}_y$ 7
 $\mathbf{T}_{xx} = \mathbf{X}'_0 \mathbf{C}\mathbf{X}_0$ 7

maximizing
 $\text{tr}(\mathbf{G}'\mathbf{A}\mathbf{G})$ subject to $\mathbf{G}'\mathbf{G} = \mathbf{I}_k$ 104
 $(\mathbf{x}'\mathbf{a})^2/\mathbf{x}'\mathbf{B}\mathbf{x}$ 106
 $(\mathbf{x}'\mathbf{A}\mathbf{y})^2/(\mathbf{x}'\mathbf{B}\mathbf{x} \cdot \mathbf{y}'\mathbf{C}\mathbf{y})$ 110
 $(\mathbf{x}'\mathbf{A}\mathbf{y})^2/(\mathbf{x}'\mathbf{x} \cdot \mathbf{y}'\mathbf{y})$ 110
 $[\mathbf{a}'(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)]^2/\mathbf{a}'\boldsymbol{\Sigma}\mathbf{a}$ 79
 $[\mathbf{a}'(\bar{\mathbf{u}} - \boldsymbol{\mu}_0)]^2/\mathbf{a}'\mathbf{S}\mathbf{a}$ 29
 $[\mathbf{a}'(\bar{\mathbf{u}}_1 - \bar{\mathbf{u}}_2)]^2/\mathbf{a}'\mathbf{S}_*\mathbf{a}$ 79
 $[\mathbf{a}'(\mathbf{u}_{(i)} - \bar{\mathbf{u}})]^2/\mathbf{a}'\mathbf{S}\mathbf{a}$ 9
 $\angle(\mathbf{V}\mathbf{z}, \mathbf{z})$ 65
 $\text{cor}(\mathbf{a}'\mathbf{x}, \mathbf{b}'\mathbf{y})$ 80
 $\text{cor}(\mathbf{y}, \mathbf{b}'\mathbf{x})$ 33
 $\text{cor}_d(\mathbf{y}, \mathbf{X}\boldsymbol{\beta})$ 20
 $\text{cor}_d(\mathbf{y}, \mathbf{X}_0\mathbf{b})$ 33
 $\cos^2(\mathbf{u}, \mathbf{v}), \mathbf{u} \in \mathcal{A}, \mathbf{v} \in \mathcal{B}$ 81
 $\frac{(\boldsymbol{\alpha}'\mathbf{A}'\mathbf{B}\boldsymbol{\beta})^2}{\boldsymbol{\alpha}'\mathbf{A}'\mathbf{A}\boldsymbol{\alpha} \cdot \boldsymbol{\beta}'\mathbf{B}'\mathbf{B}\boldsymbol{\beta}}$ 81
 $\|\mathbf{H}\mathbf{V}\mathbf{M}\|_F^2$ 66
 $\mathbf{u}'\mathbf{u}$ subject to $\mathbf{u}'\boldsymbol{\Sigma}^{-1}\mathbf{u} = c^2$ 25
 $\text{var}(\mathbf{b}'\mathbf{z})$ 78
 $\mathbf{x}'\mathbf{A}\mathbf{x}/\mathbf{x}'\mathbf{B}\mathbf{x}$ 106
 $\mathbf{x}'\mathbf{A}\mathbf{x}/\mathbf{x}'\mathbf{B}\mathbf{x}, \mathbf{B}$ nnd 108
 $\mathbf{x}'\mathbf{H}\mathbf{x}/\mathbf{x}'\mathbf{V}\mathbf{x}$ 108

- likelihood function 30
 - mean squared error 31
 - mean squared error matrix 31, 68
 - Melencolia I* by Dürer 113
 - minimizing
 - $(\mathbf{Y} - \mathbf{X}\mathbf{B})'(\mathbf{Y} - \mathbf{X}\mathbf{B})$ 22, 76, 77, 97, 117
 - $(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$ 13
 - $(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$ under $\mathbf{K}'\boldsymbol{\beta} = \mathbf{d}$ 39
 - $(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'\mathbf{V}^{-1}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$ 13, 40, 56
 - $(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'\mathbf{V}^{-1}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$ under $\mathbf{K}'\boldsymbol{\beta} = \mathbf{d}$ 41
 - $\mathbf{E}[\mathbf{y} - (\mathbf{A}\mathbf{x} + \mathbf{b})][\mathbf{y} - (\mathbf{A}\mathbf{x} + \mathbf{b})]'$ 32
 - $\text{MSEM}(\mathbf{A}\mathbf{x} + \mathbf{b}; \mathbf{y})$ 32
 - $\cos(\mathbf{V}\mathbf{z}, \mathbf{z})$ 65
 - $\text{cov}(\mathbf{G}\mathbf{y})$ subject to $\mathbf{G}\mathbf{X} = \mathbf{X}$ 52
 - $\text{cov}(\mathbf{y} - \mathbf{F}\mathbf{x})$ 32
 - $\text{cov}_d(\mathbf{Y} - \mathbf{X}\mathbf{B})$ 22, 32
 - $\text{cov}_s(\mathbf{y} - \mathbf{B}'\mathbf{x})$ 22, 32
 - $\text{eff}(\hat{\boldsymbol{\beta}}) = |\text{cov}(\hat{\boldsymbol{\beta}})|/|\text{cov}(\hat{\boldsymbol{\beta}})|$ 62
 - $\|\boldsymbol{\Sigma} - \boldsymbol{\Sigma}\mathbf{A}(\mathbf{A}'\boldsymbol{\Sigma}\mathbf{A})^{-1}\mathbf{A}'\boldsymbol{\Sigma}\|_F$ 78
 - $\|\mathbf{A} - \mathbf{B}\|_F$ subject to $\mathbf{r}(\mathbf{B}) = k$ 105, 111
 - $\|\mathbf{A} - \mathbf{B}\mathbf{Z}\|_F$ subject to \mathbf{Z} orthogonal 111
 - $\|\tilde{\mathbf{X}} - \tilde{\mathbf{X}}\mathbf{P}_G\|_F$ 78
 - $\|\mathbf{y} - \mathbf{A}\mathbf{x}\|_V$ 98
 - $\|\mathbf{y} - \mathbf{A}\mathbf{x}\|$ 95
 - $\|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|$ 13
 - $\|\mathbf{y} - \mathbf{1}\boldsymbol{\beta}\|$ 20
 - $\text{tr}(\mathbf{P}_{\boldsymbol{\Sigma}^{1/2}\mathbf{A}}\boldsymbol{\Sigma})$ 78
 - $\text{var}(\mathbf{g}'\mathbf{y})$ subject to $\mathbf{X}'\mathbf{g} = \mathbf{k}$ 52
 - $\text{var}(\mathbf{y} - \mathbf{b}'\mathbf{x})$ 33
 - $\text{var}_d(\mathbf{y} - \mathbf{X}_0\mathbf{b})$ 32, 33
 - angle 81
 - mean squared error 32
 - mean squared error matrix 32
 - orthogonal distances 78
 - minor 86
 - leading principal 86
 - principal 86
 - minus partial ordering 58, 115
 - mixed model 72
 - MLE
 - of $\sigma_{y \cdot x}^2$ 31
 - of β_0 and $\boldsymbol{\beta}_x$ 31
 - of $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ 30
 - model matrix 6
 - rank 9, 88
 - multinormal distribution
 - definition 24
 - conditional mean 26, 31
 - conditional variance 26, 31
 - contours 24
 - density function of N_2 25
 - density function of N_p 24
 - sample \mathbf{U}' from N_p 29
 - multiple correlation
 - $R = \text{cor}_d(\mathbf{y}, \hat{\mathbf{y}})$ 14
 - in \mathcal{M}_{12-1} 21
 - in no-intercept model 21
 - population 26, 31, 33
 - squared R^2 20
 - multivariate linear model 76
 - Mustonen's measure 34
- N**
- Niemelä, Jarmo vi
 - no-intercept model 5, 21, 40
 - nonnegative definiteness
 - of partitioned matrix 114
 - nonnegative matrix 112
 - norm
 - matrix 2-norm 110
 - spectral 110
 - normal distribution *see* multinormal distribution
 - normal equation 13
 - general solution 13
 - generalized 13
 - null space 9
- O**
- observation space 6
 - observation vector 6
 - OLS criterion 13
 - OLSE
 - constraints 39
 - fitted values 12
 - of $\boldsymbol{\beta}$ 14
 - of $\mathbf{K}'\boldsymbol{\beta}$ 13
 - of $\mathbf{X}\boldsymbol{\beta}$ 12, 13
 - restricted OLSE 39
 - without the i th observation 44
 - without the last b observations 46
 - ordinary least squares *see* OLSE
 - orthocomplement
 - $\mathcal{C}(\mathbf{A})^\perp$ 9
 - $\mathcal{C}(\mathbf{V}\mathbf{X}^\perp)$ 99
 - $\mathcal{C}(\mathbf{V}\mathbf{X}^\perp) = \mathcal{C}(\mathbf{W}^{-1}\mathbf{X} : \mathbf{I} - \mathbf{W}^{-1}\mathbf{W})^\perp$ 50, 99
 - $\mathcal{C}(\mathbf{V}^{-1}\mathbf{X})^\perp$ 99
 - $\mathcal{C}(\mathbf{X})_{V^{-1}}^\perp$ 99
 - \mathbf{A}^\perp 9, 84, 95
 - \mathbf{A}_V^\perp 98
 - orthogonal projector *see* projector
 - simple properties of \mathbf{P}_A 95
 - simple properties of $\mathbf{P}_{A;V}$ 98
- C** 6

H 12
J 6
M 12
 $\mathbf{P}_{22 \cdot 1}$ 97
 $\mathbf{P}_{\mathcal{C}(A) \cap \mathcal{C}(B)}$ 96
 $\mathbf{P}_{\mathcal{C}(A) \cap \mathcal{C}(B^\perp)}$ 96
 $\mathbf{P}_{\mathbf{X}; \mathbf{V}^{-1}} = \mathbf{I} - \mathbf{P}'_{\mathbf{M}; \mathbf{V}}$ 49, 99
 commuting $\mathbf{P}_A \mathbf{P}_B = \mathbf{P}_B \mathbf{P}_A$ 96
 decomposition of $\mathbf{P}_{(\mathbf{X}_1 : \mathbf{X}_2); \mathbf{V}^+}$ 99
 decomposition of $\mathbf{P}_{(\mathbf{X}_1 : \mathbf{X}_2); \mathbf{V}^{-1}}$ 57, 99
 difference $\mathbf{P}_A - \mathbf{P}_B$ 96
 sum $\mathbf{P}_A + \mathbf{P}_B$ 95
 orthogonal rotation 111
 overall-F-value 40

P

Pandora's Box
 BLUE 53, 55
 BLUP 70
 mixed model 72
 partial correlation
 $\text{pcor}_d(\mathbf{Y} | \mathbf{X})$ 21
 $\varrho_{ij \cdot \mathbf{x}}$ 26
 $\varrho_{xy \cdot z}$ 26
 $r_{xy \cdot z}$ 22
 decomposition of $1 - \varrho_{y \cdot \mathbf{x}}^2$ 34
 decomposition of $1 - R_{y \cdot 12 \dots k}^2$ 21
 decomposition of $1 - R_{y \cdot 12 \dots k}^2$ 89
 population 26
 partial covariance 26
 partitioned matrix
 g-inverse 93
 inverse 87
 MP-inverse 93
 nonnegative definiteness 114
 pencil 106, 107
 permutation
 determinant 86
 matrix 112
 Perron–Frobenius theorem 112
 Poincaré separation theorem 104
 polar decomposition 111
 predictable, unbiasedly 69
 prediction error
 $\mathbf{G}\mathbf{y} - \mathbf{y}_f$ 69
 $\mathbf{y} - \text{BLP}(\mathbf{y}; \mathbf{x})$ 26, 32
 $e_{i \cdot 1 \dots i-1}$ 34
 Mahalanobis distance 18
 with a given \mathbf{x}_* 18
 prediction interval for y_* 19
 principal component analysis 78
 matrix approximation 79
 predictive approach 78

sample principal components 78
 principal components
 from the SVD of $\tilde{\mathbf{X}}$ 79
 principal minor
 sum of all $i \times i$ principal minors 100
 principal submatrix 86
 projector *see* orthogonal projector
 $\mathbf{P}_{A|B}$ 98
 $\mathbf{P}_{\mathbf{X}|\mathbf{V}\mathbf{M}}$ 53
 oblique 95, 97
 proper eigenvalues
 $\text{ch}(\mathbf{A}, \mathbf{B})$ 107
 $\text{ch}(\mathbf{G}\mathbf{V}\mathbf{G}', \mathbf{H}\mathbf{V}\mathbf{H})$ 64
 $\text{ch}(\mathbf{V}, \mathbf{H})$ 108
 PSTricks vi

Q

QR-decomposition 111
 quadratic form
 $E(\mathbf{z}'\mathbf{A}\mathbf{z})$ 27
 $\text{var}(\mathbf{z}'\mathbf{A}\mathbf{z})$ 28
 distribution 27, 28
 independence 27, 28
 quadratic risk 68

R

random sample without replacement 23
 rank
 definition 9
 simple properties 82
 of $(\mathbf{A} : \mathbf{B})$ 83
 of $(\mathbf{1} : \mathbf{X}_0)$ 9, 88
 of $\text{cov}(\mathbf{X}\tilde{\boldsymbol{\beta}})$ 55
 of $\mathbf{A}\mathbf{B}$ 83
 of $\mathbf{A}\mathbf{P}_B$ 85
 of \mathbf{A}^- 92
 of $\mathbf{H}\mathbf{P}_V\mathbf{M}$ 64
 of $\mathbf{T}_{\mathbf{X}\mathbf{X}}$ 9, 12
 of $\mathbf{X}'(\mathbf{V} + \mathbf{X}\mathbf{U}\mathbf{X}')^{-1}\mathbf{X}$ 50
 of $\mathbf{X}'_2\mathbf{M}_1\mathbf{X}_2$ 12
 of correlation matrix 9, 88
 of model matrix 9, 88
 rank additivity
 $r(\mathbf{A} + \mathbf{B}) = r(\mathbf{A}) + r(\mathbf{B})$ 83
 Schur complement 87
 rank cancellation rule 83
 rank-subtractivity *see* minus partial ordering
 Rao's efficiency 67
 Rayleigh quotient $\mathbf{x}'\mathbf{A}\mathbf{x}/\mathbf{x}'\mathbf{x}$ 104
 recursive decomposition
 of $1 - \varrho_{y \cdot \mathbf{x}}^2$ 34
 of $1 - R_{y \cdot 12 \dots k}^2$ 21
 of $1 - R_{y \cdot 12 \dots k}^2$ 89

- of $\det(\mathbf{R})$ 89
- of $\det(\mathbf{S})$ 34
- of $\det(\mathbf{\Sigma})$ 34
- reduced model
 - $R^2(\mathcal{M}_{12\cdot 1})$ 21
 - $\mathcal{M}_{12\cdot 1}$ 61
 - \mathcal{M}_{1H} 65
 - AVP 22
 - $SSE(\mathcal{M}_{12\cdot 1})$ 21
- reducible matrix 112
- reflection 112
- regression coefficients 14
 - old ones do not change 15
 - standardized 16
- regression function 27
- regression towards mean v
- relative efficiency of OLSE *see* Watson efficiency
- residual
 - $\mathbf{y} - \text{BLP}(\mathbf{y}; \mathbf{x})$ 26, 32
 - $e_{i\cdot 1 \dots i-1}$ 34
 - after elimination of \mathbf{X}_1 20
 - externally Studentized 43, 46
 - internally Studentized 43
 - of BLUE 55, 56
 - of OLSE 13, 43
 - predicted residual 45
 - scaled 43
- residual mean square, MSE 19
- residual sum of squares, SSE 19
- rotation 111
 - of observations 79
- S**
- Samuelson's inequality 118
- Schur complement 87
 - determinant 87
 - MP-inverse 93
 - rank additivity 87
- Schur's triangularization theorem 111
- Seber, George A. F. v, vi
- shorted matrix 115
- similarity 103
- Simon, Paul v
- simultaneous diagonalization
 - by a nonsingular matrix 107
 - by an orthogonal matrix 107
 - of commuting matrices 107
- singular value decomposition 109
- skew-symmetric: $\mathbf{A}' = -\mathbf{A}$ 1
- Smith, Harry v
- solution
 - to $\mathbf{A}(\mathbf{X} : \mathbf{VX}^\perp) = (\mathbf{X}_f : \mathbf{V}_{21}\mathbf{X}^\perp)$ 70
 - to $\mathbf{AX} = \mathbf{B}$ 91
 - to $\mathbf{AXB} = \mathbf{C}$ 91
 - to $\mathbf{Ax} = \mathbf{y}$ 91
 - to \mathbf{A}^- 91
 - to $\mathbf{G}(\mathbf{X} : \mathbf{VM}) = (\mathbf{X} : \mathbf{0})$ 53
 - to $\mathbf{X}'\mathbf{X}\boldsymbol{\beta} = \mathbf{X}'\mathbf{y}$ 13
 - to $\mathbf{X}(\mathbf{A} : \mathbf{B}) = (\mathbf{0} : \mathbf{B})$ 98
- spectral radius 101, 112
- spectrum *see* eigenvalues, 100
- square root of nnd \mathbf{A} 101
- standard error of $\hat{\beta}_i$ 18
- statistical distance 9
- Stigler, Stephen M. v
- stochastic matrix 113
- stochastic restrictions 74
- sufficiency
 - linear 67
 - linear prediction 72
 - linearly minimal 68
- sum of products $\text{SP}_{xy} = t_{xy}$ 8
- sum of squares
 - change in SSE 20, 37, 39
 - change in $SSE(\mathbf{V})$ 41
 - predicted residual 45
 - $\text{SS}_y = t_{yy}$ 8
 - SSE 19
 - SSE under $\mathcal{M}_{12\cdot 1}$ 21
 - SSE, various representations 20
 - SSR 19
 - SST 19
 - weighted SSE 40, 56
- sum of squares and cubes of integers 23
- Survo vi
- Sylvester's inequality 84
- T**
- theorem
 - Albert 114
 - Cochran 28
 - Courant–Fischer 104
 - Eckart–Young 105, 111
 - Frisch–Waugh–Lovell 15, 22, 61
 - Gauss–Markov 53
 - interlacing 103
 - Lehmann–Scheffé 68
 - Perron–Frobenius 112
 - Poincaré separation 104
 - Schur's triangularization 111
 - Wedderburn–Guttman 87
- Thomas, Niels Peter vi
- Tiritiri Island 122
- total sum of squares, SST 19

transposition 86
 Trenkler, Götz vi
 triangular factorization 34
 triangular matrix 34, 111

U

unbiased estimator of σ^2
 $\hat{\sigma}^2$ 19
 $\hat{\sigma}_{(i)}^2$ 43
 $\tilde{\sigma}^2$ 40, 41, 56
 unbiasedly predictable 69

V

variable space 6
 variable vector 6
 variance
 $\text{var}_s(\mathbf{y}), \text{var}_d(\mathbf{y})$ 8
 $\text{var}(x) = E(x - \mu_x)^2$ 3
 $\text{se}^2(\hat{\beta}_i)$ 18
 $\text{var}(\mathbf{a}'\mathbf{x}) = \mathbf{a}'\boldsymbol{\Sigma}\mathbf{a}$ 10
 $\text{var}(\hat{\beta}_i)$ 17
 $\text{var}(\hat{\varepsilon}_i)$ 14
 of a dichotomous variable 23

of prediction error with a given \mathbf{x}_* 18
 variance function 27
 vec 119
 in multivariate model 76
 in multivariate sample 29
 vector of means $\bar{\mathbf{x}}$ 6
 Vehkalahti, Kimmo vi
 VIF 17, 90
 volume of the ellipsoid 38

W

Watson efficiency
 definition 62
 decomposition 66
 factorization 66
 lower limit 62
 weakly singular linear model: $\mathcal{C}(\mathbf{X}) \subset \mathcal{C}(\mathbf{V})$
 53
 Wedderburn–Guttman theorem 87
 Weisberg, Sanford v
 Wielandt inequality 117
 Wilks's lambda 30
 Wishart-distribution *see* distribution
 Working–Hotelling confidence band 19