

A model of calorimetric measurements in an open quantum system

Paolo Muratore-Ginanneschi

Department of Mathematics and Statistics University of Helsinki

In collaboration with

Brecht Donvil, Antti Kupiainen,
Jukka Pekola, Kay Schwieger.

TUNI Tampere, February 13, 2020



Outline

- 1 Calorimetric measurement on a driven qubit
- 2 Mathematical modeling
- 3 Results
- 4 Outlook


Calorimetric principle

Idea: measure work statistics in an Open Quantum System^a


^aPekola et al., “Calorimetric measurement of work in a quantum system”, 2013.

- protocols bringing the system back to the initial state at the end of the horizon.
- the work W done on the system under these conditions is equal to the heat Q dissipated to the environment

Stylized experimental setup



Integrated quantum circuit



Envisaged experimental implementation (Pekola et al., *New Journal of Physics*, (2013), Gasparinetti et al., *Physical Review Applied*, (2015), Viisanen et al., *New Journal of Physics*, (2015))

¹Schmidt, Schoelkopf, and Cleland, "Photon-Mediated Thermal Relaxation of Electrons in Nanostructures", 2004.

Closed system description

$$H = H_q + H_e + H_{qe} + H_p + H_{ep}$$

Closed system description

$$H = \textcolor{red}{H}_q + H_e + H_{qe} + H_p + H_{ep}$$

Qubit driven by a monochromatic force

$$H_q(t) = \frac{\hbar \omega_q}{2} \sigma_z + \kappa V_d(t)$$

$$V_d(t) = \hbar \omega_q (e^{i\omega_L t} \sigma_+ + e^{-i\omega_L t} \sigma_-)$$

$\kappa \hbar \omega_q$ = drive amplitude

ω_L = drive frequency

Closed system description

$$H = H_q + \textcolor{red}{H}_e + H_{qe} + H_p + H_{ep}$$

Calorimeter: free fermion gas (effectively, more follows)

$$H_e = \sum_k \eta_k c_k^\dagger c_k$$

$$\eta_k = \frac{\hbar \|\mathbf{k}\|^2}{2m}$$

Closed system description

$$H = H_q + H_e + \textcolor{red}{H}_{qe} + H_p + H_{ep}$$

Qubit calorimeter interaction

$$H_{qe} = g \frac{\sqrt{8\pi}\epsilon_F}{3N} \sum_{k \neq l \in \mathbb{S}} (\sigma_+ + \sigma_-) c_k^\dagger c_l,$$

$N = O(10^9)$ fermions

$\mathbb{S}$ = energy shell around ϵ_F

Closed system description

$$H = H_q + H_e + H_{qe} + \textcolor{red}{H}_p + H_{ep}$$

Phonons

$$H_p = \sum_k \hbar \omega_k b_k^\dagger b_k$$

$$\omega_k = v_s k$$

v_s = sound speed

$k = \|\mathbf{k}\|$ phonon wavelength norm.

Closed system description

$$H = H_q + H_e + H_{qe} + H_p + H_{ep}$$

(Herbert) Fröhlich's Hamiltonian

$$H_{ep} = \lambda \sum_{k,q} \omega_q^{1/2} \left(c_k^\dagger c_{k-q} b_q + c_k^\dagger c_{k-q} b_q^\dagger \right)$$

Timescales

- $\tau_{ee} = O(10^0)\text{ns}$:
Landau quasi-particle relaxation rate to Fermi–Dirac equilibrium in a metallic wire.
- $\tau_{ep} = O(10^4)\text{ns}$:
electron-phonon interactions.
- $\tau_R = 2 - 5 \times O(10^5)\text{ns}$: transmon qubit relaxation times (Wang et al., *Applied Physics Letters*, (2015))
- $\tau_{eq} \simeq g^{-2}$
Fermi's golden rule estimate of characteristic **qubit-calorimeter time scale**.

Open quantum system approach

$$\tau_{ee} \ll \tau_{eq} \ll \tau_{ep} \ll \tau_R$$



Phonon–fermion bath interaction

- Phonon bath temperature $T_p = O(10^{-1})\text{K}$ (cryostat)
- Fermion bath temperature T_e

$$T_e \simeq T_p$$

mean energy current $\propto T_p^5 - T_e^5$ (leading order^a)

rms energy current fluctuations $\propto O(T_p^3)$ at $T_e = T_p$ (leading order^b)

^aKaganov, Lifshitz, and Tanatarov, “Relaxation between Electrons and the Crystalline Lattice”, 1957; Wellstood, Urbina, and Clarke, “Hot-electron effects in metals”, 1994.

^bPekola and Karimi, “Quantum noise of electron-phonon heat current”, 2018.

Idea of the model

Qubit: stochastic Schrödinger equation^a

^aBreuer and Petruccione, *The Theory of Open Quantum Systems*, 2002.

$$d\psi = (\text{deterministic dissipative drift}) dt + \text{Poisson jumps}$$

Calorimeter: equilibrium Fermi–Dirac ensembles at evolving T_e ^a

^aBerg, Brange, and Samuelsson, “Energy and temperature fluctuations in the single electron box”, 2015; Marinari and Parisi, “Simulated Tempering: A New Monte Carlo Scheme”, 1992.

$$dT_e^2 = \frac{1}{N\gamma} dE$$

Sommerfeld expansion

$$dE = dE_{eq} + dE_{ep} = \text{Poisson jumps} + (T_p^5 - T_e^5)dt + O(T_p^3)dw_t$$



Stochastic Jump Process

Closed system: unitary evolution

$$\psi(t + dt) - \psi(t) = d\psi(t) = -iH\psi dt$$

Open quantum system: stochastic Schrödinger equation²³⁴

- Fermi golden rule: **dissipative terms** are added to the Hamiltonian

$$H\psi(t)dt \rightarrow G(\psi(t))dt$$

- Fermi golden rule: transitions induce **stochastic jumps**

$$(|\pm\rangle - \psi(t)) dN(\mp\omega), \quad dN(\mp\omega) = 0, 1,$$

$$\mathbb{E}_\psi(dN(\mp\omega)) = \gamma(\mp\omega) \|A(\mp\omega)\psi\|^2 dt$$

Weak-drive approach: add drive as a perturbation to the continuous evolution

$$G(\psi(t)) + \kappa H_d(t)\psi(t)$$

²Ghirardi, Rimini, and Weber, “Unified dynamics for microscopic and macroscopic systems”, 1986.

³Dalibard, Castin, and Mølmer, “Wave-function approach to dissipative processes in quantum optics”, 1992.

⁴Breuer and Petruccione, *The Theory of Open Quantum Systems*, 2002.

Temperature Process


Using the Sommerfeld expansion we find the dependence of the temperature on the change in internal energy E of the calorimeter

$$dT_e^2 = \frac{dE_{eq} + dE_{ep}}{\gamma}.$$

The qubit-electron interaction **alone** ($dE_{ep} = 0$) gives

$$dE_{eq} = \hbar\omega(dN(\omega) - dN(-\omega)),$$

$$d\psi(t) = -i[G(\psi(t)) + \kappa H_d(t)\psi(t)]dt \\ + \left(|+\rangle - \psi(t)\right)dN(-\omega) + \left(|-\rangle - \psi(t)\right)dN(\omega)$$



Upshot of the modeling

"Strong drive": Floquet theory^a

^aBreuer and Petruccione, "Dissipative quantum systems in strong laser fields: Stochastic wave-function method and Floquet theory", 1997.

- $\tau_{qe} \gg \tau_m$ = inverse separation of peaks in the radiation spectrum (RWA).
- Resonant drive: $\tau_m/\tau_{qe} \simeq g^2/\kappa \ll 1$
- Temperature+population process: **jump diffusion master equation**

"Weak drive"


- $g^2/\kappa \geq 1$
- Temperature+state process: **hybrid master equation^a**

^aChruściński et al., "Dynamics of Interacting Classical and Quantum Systems", 2011.



Short-time temperature behaviour



Initial temperature of the electron bath: $T_e = 0.1K$



Temperature distributions after 10 periods of resonant **strong drive**

Short-time temperature behaviour

Initial temperature of the electron bath: $T_e = 0.1K$




(Left) Mean temperature of the calorimeter after 10 periods of driving vs driving frequency ω_L for different values of the qubit calorimeter coupling g .


Stars= weak-drive. Lines: Floquet . (Right) Standard deviation.

Short-time temperature behaviour

Initial temperature of the electron bath: $T_e = 0.1K$



Relaxation to a steady state



The qubit-calorimeter reaches a steady state.

Effective temperature process

Multiscale expansion: $\varepsilon \propto 1/N$ & $s = \varepsilon t \geq O(1)$

$$dT_e^2 = \frac{1}{\gamma} \left(\Sigma V (T_p^5 - T_e^5) + J(T_e^2) \right) ds + \frac{1}{\gamma \sqrt{N}} \left(\sqrt{10 \Sigma V k_B} T_p^3 + \sqrt{S(T_e^2)} \right) dw_s$$


Analytic estimates

$$\langle T_e \rangle \approx \left(T_p^5 + g^2 \frac{O(\hbar \omega_L^2)}{\Sigma V} \right)^{1/5} \quad \text{mean steady state temperature}$$

$$\tau \approx \left(T_p^5 + g^2 \frac{O(\hbar \omega_L^2)}{\Sigma V} \right)^{-3/5} \quad \text{relaxation time to steady state}$$




Numerics vs analytic theory (Floquet)



Steady state temperature PDF

Steady state av. temperature vs drive strength



Outlook

- Predictions always involve weak coupling between qubit and calorimeter.
- Perturbative Markovian master equation techniques not reliable beyond the strictly weak subsystem-bath coupling limit (see e.g. Segal, *Physical Review B*, (2013)).
- Strong qubit-calorimeter coupling analysis desirable.

References

Based on

- A. KUPIAINEN, P. MURATORE-GINANNESCHI, J.P. PEKOLA, and K. SCHWIEGER *Fluctuation Relation for Qubit-Calorimetry* arXiv:1606.02984, and Phys. Rev. E. 94, 062127, (2016).
- B. DONVIL, P. MURATORE-GINANNESCHI, J. P. PEKOLA, and K. SCHWIEGER, *A model for calorimetric measurements in an open quantum system* arXiv:1803.11015 and Phys. Rev. A 97, 052107 (2018).
- B. DONVIL, P. MURATORE-GINANNESCHI, J. P. PEKOLA *Hybrid master equation for calorimetric measurements* arXiv:1811.12832 and Phys. Rev. A, 99, 042127 (2019)

THANKS FOR YOUR ATTENTION



THANKS, Brecht, Antti, Jukka & Kay

